



PROTECTIVE EFFECT OF PHYTOCHEMICAL CONSTITUENTS ON OXIDATIVE & NON OXIDATIVE DNA DAMAGE INDUCED BY H₂O₂ & ASPIRIN

Dr. Eswari Beeram^{1*}, P Govardhan^{1a}, Dr. S. Hemachandra², S. Himagirish Kumar^{2a},
Swetha Madamala³,

Dr. B. Sambasivaiah^{2b}, Kartheek Rao Amperayani^{3a}

Assistant Professor, DBCS, SOLAS, Mohan Babu University, A.P., India.

Master of Biotechnology, DBCS, SOLAS, Mohan babu University, A.P., India.

Principal, Sanskrithi School of Engineering, Puttaparthi, A.P., India.

Assistant Professor, Dept. Of Organic Chemistry, Sri Padmavathi Mahila Viswavidyalayam,
A.P., India.

Assistant Professor, Department of Chemistry, MVJ College of Engineering, Karnataka,
India.

Associate Professor, Department of Chemistry, Sanskrithi School of Engineering, Puttaparthi,
A.P., India.

Assistant Professor, Dept. Of Chemistry, Gayatri Vidya Parishad College of Degree and PG
courses, Rushikonda, Visakhapatnam, A.P., India.

Corresponding author Email: eshu.sonu@gmail.com, Mobile:9700277136.

Abstract

Aspirin is a non steroidal anti inflammatory drug used widely due to its low cost and efficacy. Aspirin alone does not cause DNA damage but at certain situations like aspirin in water can induce DNA damage. H₂O₂ is generally used to induce oxidative stress and in this study we used 1% Aspirin dissolved in water and three different concentrations of H₂O₂ like 5%,15% and 25% to induce DNA damage. We studied protective effect of 10 phytochemical constituents like *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus* on DNA damage induced by aspirin and H₂O₂. In the 10 phytochemicals used *Oryza sativa* offered full protection commonly to all three concentrations of 5%,15% and 25% H₂O₂ used. Where as *Lablab purpureus* *Ocimum sativum* and *Eucalyptus* protected commonly with 5% and 25% H₂O₂.

Key words: Aspirin, H₂O₂, Oxidative stress, NSAIDs, *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus*

Introduction:

Aspirin is a non steroidal anti inflammatory drug used widely due to its low cost and efficacy. From the studies of Gómez-Oliván LM et al.,(2014) aspirin dissolved in water solution cause oxidative stress and DNA damage in *Daphnia magna*. Inflammation is one of the major cause for bone marrow ablation induced after irradiation and from the studies of Dorota Kacprzak , Rafał

Pawliczak (2015). aspirin protects bone marrow ablation by reducing the antiinflammatory genes gene expression.

Aspirin induced asthma is a distinct disorder caused by ingestion of aspirin or other anti-inflammatory non steroidal drugs (NSAIDs). However more recent studies are carried on the study of oxidative stress induced by aspirin in the cancer cells. However the patho mechanism is not well understood (Dorota Kacprzak , Rafał Pawliczak (2015)) . From the studies of Gąsecka et al., (2025) it is proven to be oxidative stress induced due to coronary artery bypass grafting is independent of aspirin intake.

Generally H₂O₂ is the preferred agent to induce oxidative stress and according to the reports of Ransy C et al., (2020) high concentrations of H₂O₂ is required to induce oxidative stress and can be monitored using polarographic measurement and mechanism of oxidative stress by H₂O₂ involves production of O₂ and exposure of O₂ to uncoupled NOS and NADPH oxidase. According to findings of Ransy C et al., (2020) complex interaction between H₂O₂ and oxygen Radical generating enzymes is responsible for the oxidative stress induced by H₂O₂. In this current study we studied the effect of DNA damage induced by Aspirin and H₂O₂ and protection offered by various phytochemical constituents against DNA damage induced by aspirin and H₂O₂.

Methodology:

1. Isolation of DNA from chicken liver:

DNA is isolated from chicken liver using the procedure of Vijay Hemmadi (2016).

2. Separation of DNA by Agarose Gel Electrophoresis:

Seperation of DNA by agarose gel electrophoresis was done according to vlab.amritha.edu., (2012). Agarose Gel Electrophoresis (AGE).

3. Induction of DNA damage by aspirin and 5%,15% and 25% H₂O₂:

1ml of isolated DNA is transferred to standard flask and made up to 100ml with distilled water and used as standard for studying the DNA damage. 1% aspirin solution is prepared by weighing and transferring of 1g of aspirin in to 100ml distilled water. Similarly 30% H₂O₂ is used for preparing stock solutions of 5%, 15% and 25% H₂O₂ and these stock solutions are used for inducing oxidative damage in DNA.

To 1ml of DNA solution, 1ml of aspirin and 5%,15% and 25% H₂O₂ is added to different test tubes and incubated for time periods 15', 30',45' and 60' time period and the reading is recorded at 260nm using UV Visible spectrophotometer.

4. Study of protective effect of phytochemical on DNA damage induced by Aspirin and 5%,15% and 25% H₂O₂:

1g of phytochemical is dissolved in 10ml of distilled water and allowed to stand for 1 hr. After the time period 1ml of the filtrate of phytochemical is added to tubes with aspirin + DNA and similarly to DNA+5%,15% and 25% H₂O₂ separately. The reaction is stopped by addition of 0.5ml of 0.5M NaOH. Reading is recorded after 15', 30',45' and 60' time period using UV Visible spectrophotometer.

The results are tabulated and represented graphically to study the protective effect of phytochemicals on DNA damage.

10 phytochemical constituents like *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus* are used for the study.

Calculation of molarity of H₂O₂:

5% H₂O₂: 5ml of 30% H₂O₂ is added to 100ml of water and the molarity of 5% H₂O₂ is calculated to be as 0.44M.

15% H₂O₂: 15ml of 30% H₂O₂ is added to 100ml of water and the molarity of 5% H₂O₂ is calculated to be as 1.32M.

25% H₂O₂: 25ml of 30% H₂O₂ is added to 100ml of water and the molarity of 5% H₂O₂ is calculated to be as 2.2M.

Results:

DNA is isolated from chicken liver and separated on 1% agarose gel. Oxidative and non oxidative damage of DNA is studied by inducing DNA damage with aspirin and H₂O₂. Protective effect of phytochemicals against DNA damage is studied using Phyto constituents like *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus*.





Figure:1 Separation of DNA on 1% agarose gel.

Oxidative damage is studied using concentrations of 5%,15% and 25% H₂O₂ to induce oxidative stress and from table 1 after 15' of exposure phytochemicals except *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca* and *Vigna radiata* offered protection against DNA damage but after 60min exposure *Vigna mungo* failed to offer protection against DNA damage induced by 5% H₂O₂..

	5% H ₂ O ₂ + DNA	Lab lab	Man gifer a indic a	Vigna ungui culata	Mus a	Oryz a sativ a	Vigna radia ta	Vign a mun go	Aza dira chta	Ocim um	Euca lyptu s
15'	10	- 0.04 21	10	10	10	- 1.262 7	10	- 0.507 9	0.17 23	- 0.527 9	- 2.297 6
30'	- 0.048 9	0.34 4	10	0.847 5	10	- 0.663 1	0.571 1	10	0.54 57	10	- 0.870 2
45'	- 0.265 3	- 0.07 41	10	10	10	- 0.698 8	1.128 2	10	10	- 0.926 5	- 0.168 3
60'	- 0.023 3	- 0.12 57	10	0.521 7	10	- 0.595 7	- 0.009 3	10	- 0.30 63	- 0.763 1	- 0.916 2

Table: 1 Effect of 5% H₂O₂ on DNA and protection against DNA damage offered by phytochemical constituents *Lablab purpureus*, *Oryza sativa*, *Vigna radiata*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus*.



Graph: 1 Protection offered by Phytochemical constituents against DNA damage induced by 5% H2O2.


	15% H2O2 + DNA	Lablab	Mangifera indica	Vigna unguiculata	Musa	Oryza sativa	Vigna radiata	Vigna mungo	Azadirachta	Ocimum	Eucalyptus
15'	-0.8951	0.0038	-0.1653	1.4456	0.2858	0.1226	-1.0714	0.9848	10	-0.228	10
30'	-0.2962	1.0117	-0.0619	10	0.5462	-0.1123	10	0.1736	2.4774	10	-0.076
45'	-0.8195	10	10	10	0.6695	-0.2323	10	10	3.191	10	10
60'	1.2215	10	-0.0559	2.1119	-0.0051	-0.0644	10	10	10	10	10

Table:2 Effect of 15% H2O2 on DNA and protective effect of *Vigna radiata*, *Mangifera indica*, *Musa paradisiaca* and *Oryza sativa* on DNA damage.

15' min exposure of DNA to 15% H2O2 resulted in DNA damage and in all the 10 phytochemicals used only *Vigna radiata* offered protection against DNA damage for 15' exposure and exposure to 60' resulted in increase in DNA damage compared to 15' exposure and phytoconstituents like *Mangifera indica*, *Musa paradisiaca* and *Oryza sativa* offered protection against DNA damage induced by 15% H2O2.

25% H2O2 induced greater DNA damage compared to 15% H2O2 and 15' min exposure resulted in highest percentage of DNA damage and almost all phytochemicals except *Vigna radiata*, *Vigna mungo* and *Azadirachta indica* offered protection against DNA damage induced by 25% H2O2. Decrease in absorbance after 60min exposure is may be due to reannealing of the repetitive DNA


and phytochemicals like *Mangifera indica* elevated the DNA damage induced by 25% H₂O₂. *Vigna mungo* offered protection against DNA damage after 60' exposure to 25% H₂O₂ may be due to DNA aggregation.



Graph: 2 Protective effect of phytochemicals *Vigna radiata*, *Mangifera indica*, *Musa paradisiaca* and *Oryza sativa* on DNA damage induced by 15% H₂O₂

	25% H ₂ O ₂ + DNA	Lablab	Mangifera indica	Vigna unguiculata	Musa	Oryza sativa	Vigna radiata	Vigna mungo	Azadirachta	Ocimum	Eucalyptus
15'	10	- 0.43 05	0.335 8	- 0.334 8	- 0.382 1	0.12 03	10	10	10	- 0.242 6	0.689 4
30'	0.054 8	- 0.04 31	- 0.079 3	- 0.817 8	- 0.803 1	- 0.65 51	10	0.432 9	10	- 0.455 3	0.061 4
45'	0.297 4	- 0.48 33	- 0.088 8	10	- 0.569	- 0.78 94	- 0.517 8	0.383	0.240 2	-0.717	0.194 2
60'	0.839 1	0.56 98	10	- 0.275 6	- 0.683	- 0.23 77	10	0.382 5	10	-0.543	- 0.256 8


Table: 3 Protective effect of phytochemicals *Lablab purpureus*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Ocimum sativum* and *Eucalyptus* on DNA damage induced by 25% H₂O₂.



Graph: 3 Protection offered against DNA damage by Phyto constituents *Lablab purpureus*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Ocimum sativum* and *Eucalyptus*.

	Aspirin + DNA	<i>Lablab</i>	<i>Mangifera indica</i>	<i>Vigna unguiculata</i>	<i>Musa</i>	<i>Oryza sativa</i>	<i>Vigna radiata</i>	<i>Vigna mungo</i>	<i>Azadirachta</i>	<i>Ocimum</i>	<i>Eucalyptus</i>
15'	10	-0.1943	10	-0.7688	0.2166	-0.2484	0.3629	10	10	0.7542	0.0077
30'	-0.3242	10	-0.3242	-0.825	0.3387	10	10	10	10	0.2498	0.6634
45'	0.4099	-0.438	0.4099	-0.6337	10	0.1898	10	10	10	0.6182	0.3514
60'	-0.373	-0.4482	-0.373	-0.4039	10	10	10	10	10	0.1374	0.2265

Table: 4 Protection against oxidative and non oxidative DNA damage induced by Aspirin by Phytochemical constituents like *Lablab purpureus*, *Vigna unguiculata* after 60' exposure.




Graph: 4: Bar graph on protective effect of phyto constituents *Lablab purpureus*, *Vigna unguiculata* on DNA damage induced by Aspirin


Lablab purpureus, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Ocimum sativum* and *Eucalyptus* offered protection against DNA damage after 15' and 60' min exposure of DNA to the drug aspirin, but *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Ocimum sativum* and *Eucalyptus* failed to protect DNA against damage by aspirin after 60 'min exposure.

From the graph 5 the DNA damage caused by aspirin is found to be about 23% and most of the phytochemicals like *Lablab purpureus*, *Vigna unguiculata* showed protective effect with percentage of 81% and 99%.


From the graph 6 DNA damage induced by 5% H₂O₂, 15% H₂O₂ and 25% H₂O₂ is found to be 18%, -5% and 23%.




Graph: 5: % of DNA protection offered by various phytochemicals after treatment with Aspirin.



Graph: 6 Percentage of protection against DNA Damage by Phytochemicals constituents after treatment with 5% H₂O₂



Graph:7 Percentage of protective effect offered against DNA damage induced by 15% H₂O₂ by various phytochemicals like *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus*.



Graph: 8: Protection percentage of DNA offered by phytochemicals like *Lablab purpureus*, *Mangifera indica*, *Vigna unguiculata*, *Musa paradisiaca*, *Oryza sativa*, *Vigna radiata*, *Vigna mungo*, *Azadirachta indica*, *Ocimum sativum* and *Eucalyptus* against DNA damage induced by 25% H₂O₂.

Discussion and Conclusion:

H₂O₂ is one of the key inducer of oxidative stress mediated intracellularly through redox signaling leading to loss of cellular integrity. From the findings of Saputra et al., (2024) H₂O₂ induced oxidative stress in zebra fish causes impairment of embryonic development and hatching and can be negotiated by the addition of antioxidants like GSH. H₂O₂ exposure can reduce the cell viability in meningotheial cells and from the studies of Xin et al.,(2022) H₂O₂ exposure affects meningotheial cells by increasing oxidative stress, decreasing antioxidant enzymes, increasing level of free radicals and decrease in mitochondrial membrane potential compared to unexposed cells and responsible for toxicity in the cells.

Aspirin can induce oxidative stress and DNA damage at certain conditions when the exposure is for greater period. From the findings of Jiang et al.,(2023) inhibition of glutathione synthesis increases the cytotoxic effect of aspirin and however exposure to N-acetyl cysteine can augments the cytotoxicity caused (Haider Raza,Annie John (2012)). From the current study oxidative stress caused by H₂O₂ and the oxidative and non oxidative DNA damage induced by aspirin can be augmented by certain phytoactive components *Oryza sativa*,*Lablab purpureus* *Ocimum sativum* and *Eucalyptus* used in the study.

References:

1. Dorota Kacprzak , Rafał Pawliczak (2015). Does aspirin-induced oxidative stress cause asthma exacerbation?Arch Med Sci;494–504. doi: [10.5114/aoms.2014.41960](https://doi.org/10.5114/aoms.2014.41960).
2. Gąsecka et al., (2025). Effect of aspirin dosage on oxidative stress and platelet reactivity in patients undergoing coronary artery bypass grafting (APRICOT): randomized controlled trial.Platelets Volume 36,Issue 1 <https://doi.org/10.1080/09537104.2025.2457415>
3. Gómez-Oliván LM et al.,(2014).DNA damage and oxidative stress induced by acetylsalicylic acid in *Daphnia magna*.Comp Biochem Physiol C Toxicol Pharmacol; Vol.164:21-6.doi: [10.1016/j.cbpc.2014.04.004](https://doi.org/10.1016/j.cbpc.2014.04.004). Epub
4. [Jiang et al.,\(2023\).Aspirin protects against genotoxicity by promoting genome repair; Cell Research volume 33, pages 325–327.](#)
5. Ransy C et al., (2020). Use of H₂O₂ to Cause Oxidative Stress, the Catalase Issue. Int J Mol Sci;30;21(23):9149. doi: [10.3390/ijms21239149](https://doi.org/10.3390/ijms21239149); PMID: [33266350](https://pubmed.ncbi.nlm.nih.gov/33266350/) PMCID: [PMC7731207](https://pubmed.ncbi.nlm.nih.gov/PMC7731207/)
6. [Saputra et al., \(2024\).Oxidative stress induced by hydrogen peroxide disrupts zebra fish visual development by altering apoptosis, antioxidant and estrogen related genes; Scientific Reports volume 14, Article number: 14454..](#)
7. [Xin et al.,\(2022\). Hydrogen peroxide initiates oxidative stress and proteomic alterations in meningotheial cells; Scientific Reports volume 12, Article number: 14519.](#)

8. HaiderRaza,AnnieJohn (2012). Implications of Altered Glutathione Metabolism in Aspirin-Induced Oxidative Stress and Mitochondrial Dysfunction in HepG2 Cells. PLoSONE;Volume7,Issue4,e36325.
9. Vijay Hemmadi (2016). Extraction of DNA from chicken liver. Method; IMET; DOI : 10.13140/RG.2.1.2807.5924.
10. Seperation of DNA by agarose gel electrophoresis. vlab.amritha.edu,. (2012). Agarose Gel Electrophoresis (AGE).